arXiv:1406.0378v1 [math.CO] 2 Jun 2014

Connective eccentric index of some graph operations

Nilanjan De[a], Anita Pal[b], Sk. Md. Abu Nayeem[c,*]

[a]*Department of Basic Sciences and Humanities (Mathematics), Calcutta Institute of Engineering and Management, Kolkata, India.*
[b]*Department of Mathematics, National Institute of Technology, Durgapur, India.*
[c]*Department of Mathematics, Aliah University, DN - 20, Sector - V, Salt Lake, Kolkata - 700 091, India.*

Abstract

The connective eccentric index of a graph is a topological index involving degrees and eccentricities of vertices of the graph. In this paper, we have studied the connective eccentric index for double graph and double cover. Also we give the connective eccentric index for some graph operations such as joins, symmetric difference, disjunction and splice of graphs.

MSC (2010): Primary: 05C35; Secondary: 05C07, 05C40

Keywords: Eccentricity, degree, graph invariant, connective eccentric index, graph operations.

1. Introduction

Let G be a simple connected graph with vertex set $V(G)$ and edge set $E(G)$. Let n and m be the number of vertices and edges of G respectively. We denote the degree of a vertex v of G by $d_G(v)$. For $u, v \in V(G)$, the distance between u and v is defined as the length of a shortest path between u and v. For a given vertex v of G, the eccentricity $\varepsilon_G(v)$ is the largest distance from v to any other vertices of G. The sum of eccentricities of all the vertices of G is denoted by $\theta(G)$ [3]. If any vertex $v \in V(G)$ is adjacent to all the other vertices of G then v is called a well-connected vertex. Thus, if $v \in V(G)$ is a well-connected vertex, then $\varepsilon_G(v) = 1$. For example, all the vertices of a complete graph are well-connected.

Recently, a number of topological indices involving vertex degree and eccentricity were subject to a lot of mathematical as well as chemical studies. A topological index of this type, introduced by Gupta et al. [10], was named as the connective eccentric index and was defined as

$$C^{\xi}(G) = \sum_{v \in V(G)} \frac{d_G(v)}{\varepsilon_G(v)}.$$

*Corresponding Author.
Email addresses: de.nilanjan@rediffmail.com (Nilanjan De), anita.buie@gmail.com (Anita Pal), nayeem.math@aliah.ac.in (Sk. Md. Abu Nayeem)

M. Ghorbani [7] gave some bounds of connective eccentricity index and also computed this index for two infinite classes of dendrimers. De [4] reported some bounds for this index in terms of some graph invariants such as maximum and minimum degree, radius, diameter, first Zagreb index and first Zagreb eccentricity index, etc. In [8], Ghorbani and Malekjani computed the eccentric connectivity index and the connective eccentric index of an infinite family of fullerenes. In [17], Yu and Feng also derived some upper or lower bounds for the connective eccentric index and investigated the maximal and the minimal values of connective eccentricity index among all n-vertex graphs with fixed number of pendent vertices.

In [7], Ghorbani showed that for a vertex transitive graph G, the connective eccentric index is given by

$$C^{\xi}(G) = \sum_{j=1}^{t} |A_j| \frac{d_G(x_i)}{\varepsilon_G(x_i)},$$

where $A_1, A_2, \ldots, A_t$ are the orbits of $Aut(G)$ under its natural action on $V(G)$ and $x_i \in A_i, 1 \leq i \leq t$. In particular, if G is a regular graph, then $C^{\xi}(G) = \dfrac{n\delta}{r(G)}$, where n is the number of vertices of G, which is a δ-regular graph and $r(G)$ is the radius of G. Let, $K_n, C_n, Q_m, \Pi_m, A_m$ denote the complete graph with n vertices, the cycle on n vertices, m-dimensional hypercube, m-sided prism and the m-sided antiprism respectively. It can be easily verified that the explicit formulae for the connective eccentric index of $K_n, C_n, Q_m, \Pi_m, A_m$ are as follows.

Proposition 1. $C^{\xi}(K_n) = n(n-1)$.

Proposition 2. $C^{\xi}(C_n) = \frac{2n}{\lfloor n/2 \rfloor}$.

Proposition 3. $C^{\xi}(Q_m) = 2^m$.

Proposition 4. $C^{\xi}(\Pi_m) = \begin{cases} \frac{12m}{m+2}, & \textit{when } m \textit{ is even} \\ \frac{12m}{m+1}, & \textit{when } m \textit{ is odd.} \end{cases}$

Proposition 5. $C^{\xi}(A_m) = \begin{cases} 16, & \textit{when } m \textit{ is even} \\ \frac{16m}{m+1}, & \textit{when } m \textit{ is odd.} \end{cases}$

Several studies on different topological indices related to graph operations of different kinds are available in the literature [2, 9, 11, 14, 15].

In this paper, first we calculate connective eccentric index of double graph and double cover and hence the explicit formulae for the connective eccentric indices of join, symmetric difference, disjunction and splice of graphs are obtained. For the definitions and different results on graph operations, such as join, symmetric difference, disjunction, etc., readers are referred to the book of Imrich and Klavžar [13].

2. Main Results

In this section, first we define and then compute eccentric connectivity index of double graph and double cover graph.

2.1. Connective eccentric index of double graph and double cover

Let us denote the double graph of a graph G by G^*, which is constructed from two copies of G in the following manner [1, 12]. Let the vertex set of G be $V(G) = \{v_1, v_2, \ldots, v_n\}$, and the vertices of G^* are given by the two sets $X = \{x_1, x_2, \ldots, x_n\}$ and $Y = \{y_1, y_2, \ldots, y_n\}$. Thus for each vertex $v_i \in V(G)$, there are two vertices x_i and y_i in $V(G^*)$. The double graph G^* includes the initial edge set of each copies of G, and for any edge $v_iv_j \in E(G)$, two more edges x_iy_j and x_jy_i are added.

Figure 1: The graph P_3 and its double graph P_3^*.

Theorem 1. *The connective eccentric index of the double graph G^* is given by $C^{\xi}(G^*) = 4C^{\xi}(G) - 2(n-1)\|n-1\|_G$, where, $\|n-1\|_G$ is the number of vertices with degree $n-1$, i.e., of eccentricity one.*

Proof. From the construction of double graph, it is clear that $d_{G^*}(x_i) = d_{G^*}(y_i) = 2d_G(v_i)$, where $v_i \in V(G)$ and $x_i, y_i \in V(G^*)$ are the corresponding clone vertices of v_i. Also we can write $\varepsilon_{G^*}(x_i) = \varepsilon_{G^*}(y_i) = \varepsilon_G(v_i)$, when $\varepsilon_G(v_i) \geq 2$ and $\varepsilon_{G^*}(x_i) = \varepsilon_{G^*}(y_i) = \varepsilon_G(v_i) + 1$ when $\varepsilon_G(v_i) = 1$.

Thus the connective eccentric index of double graph G^* is

$$\begin{aligned}
C^{\xi}(G^*) &= \sum_{i=1}^{n} \frac{d_{G^*}(x_i)}{\varepsilon_{G^*}(x_i)} + \sum_{i=1}^{n} \frac{d_{G^*}(y_i)}{\varepsilon_{G^*}(y_i)} \\
&= 2\left[\sum_{\varepsilon_G(v_i)\geq 2} \frac{2d_G(v_i)}{\varepsilon_G(v_i)} + \sum_{\varepsilon_G(v_i)=1} \frac{2d_G(v_i)}{\varepsilon_G(v_i)+1}\right] \\
&= 4\left[\sum_{\varepsilon_G(v_i)\geq 2} \frac{d_G(v_i)}{\varepsilon_G(v_i)} + \sum_{\varepsilon_G(v_i)=1} \frac{d_G(v_i)}{2}\right] \\
&= 4\left[\sum_{\varepsilon_G(v_i)\geq 2} \frac{d_G(v_i)}{\varepsilon_G(v_i)} + \sum_{\varepsilon_G(v_i)=1} \frac{d_G(v_i)}{1}\right] - 2\sum_{\varepsilon_G(v_i)=1} d_G(v_i) \\
&= 4C^{\xi}(G) - 2(n-1)\|n-1\|_G.
\end{aligned}$$

□

Let $G = (V, E)$ be a simple connected graph with $V = \{v_1, v_2, ..., v_n\}$. The extended double cover of G, denoted by G^{**} is the bipartite graph with bipartition (X, Y) where $X = \{x_1, x_2, ..., x_n\}$ and $Y = \{y_1, y_2, ..., y_n\}$ in which x_i and y_j are adjacent if and only if

either v_i and v_j are adjacent in G or $i = j$. For example, the extended double cover of the complete graph is the complete bipartite graph. This construction of the extended double cover was introduced by Alon [1] in 1986.

Figure 2: The graph P_3 and its extended double cover P_3^{**}.

Theorem 2. *The connective eccentric index of the extended double cover G^{**} satisfies the inequality*

$$C^{\xi}(G^{**}) \leq 2C^{\xi}(G) - (n-2)\|n-1\|_G.$$

Proof. If G is a graph with n vertices and m edges then, from definition, the extended double cover graph G^{**} consists of $2n$ vertices and $(n+2m)$ edges and $d_{G^{**}}(x_i) = d_{G^{**}}(y_i) = d_G(v_i) + 1$ and $\varepsilon_{G^*}(x_i) = \varepsilon_{G^*}(y_i) = \varepsilon_G(v_i) + 1$ for $i = 1, 2, \ldots, n$.

Thus the connective eccentric index of extended double cover graph G^{**} is given by

$$\begin{aligned}
C^{\xi}(G^{**}) &= \sum_{i=1}^{n} \frac{d_{G^{**}}(x_i)}{\varepsilon_{G^{**}}(x_i)} + \sum_{i=1}^{n} \frac{d_{G^*}(y_i)}{\varepsilon_{G^*}(y_i)} \\
&= 2\left[\sum_{\varepsilon_G(v_i)\geq 2} \frac{d_G(v_i)+1}{\varepsilon_G(v_i)+1} + \sum_{\varepsilon_G(v_i)=1} \frac{d_G(v_i)+1}{2}\right] \\
&\leq 2\sum_{\varepsilon_G(v_i)\geq 1} \frac{d_G(v_i)}{\varepsilon_G(v_i)} + \sum_{\varepsilon_G(v_i)=1} \{d_G(v_i)+1\} \\
&= 2\left[\sum_{\varepsilon_G(v_i)\geq 1} \frac{d_G(v_i)}{\varepsilon_G(v_i)} + \sum_{\varepsilon_G(v_i)=1} d_G(v_i)\right] - \sum_{\varepsilon_G(v_i)=1} d_G(v_i) + \sum_{\varepsilon_G(v_i)=1} 1 \\
&= 2C^{\xi}(G) - (n-1)\|n-1\|_G + \|n-1\|_G \\
&= 2C^{\xi}(G) - (n-2)\|n-1\|_G.
\end{aligned}$$

□

Now some exact formula for the eccentric connectivity index of joins, symmetric difference, disjunction, splice graphs are presented.

2.2. Join

The join $G_1 + G_2$ of two graphs G_1 and G_2 with disjoint vertex sets V_1, V_2 and edge sets E_1, E_2, is the graph union $G_1 \cup G_2$ together with all the edges joining V_1 and V_2, i.e., $G_1 + G_2$ consists of the vertex set $V_1 \cup V_2$ and edge set $E_1 \cup E_2 \cup \{xy : x \in V_1, y \in V_2\}$.

Theorem 3. *Let G_1 and G_2 be two graphs without well-connected vertices. Then, $C^{\xi}(G_1 + G_2) = |E(G_1)| + |E(G_2)| + |V(G_1)||V(G_2)|$.*

Proof. Let n_1 and n_2 be the numbers of vertices in G_1 and G_2 respectively. Thus, $|V(G_1 + G_2)| = |V_1| + |V_2| = n_1 + n_2$. For vertices $u \in V_1$ and $v \in V_2$, it holds that $d_{G_1+G_2}(u) = d_{G_1}(u) + n_2$ and $d_{G_1+G_2}(v) = d_{G_2}(v) + n_1$. Since none of G_1 and G_2 contains well connected vertices, then for every $u \in V(G_1 + G_2), \varepsilon_{G_1+G_2}(u) = 2$. So, from definition of connective eccentric index, we have

$$\begin{aligned} C^{\xi}(G_1 + G_2) &= \sum_{v\in V(G_1)} \frac{d_{G_1+G_2}(v)}{\varepsilon_{G_1+G_2}(v)} + \sum_{v\in V(G_2)} \frac{d_{G_1+G_2}(v)}{\varepsilon_{G_1+G_2}(v)} \\ &= \frac{1}{2}\left[\sum_{v\in V(G_1)} \{d_{G_1}(v) + n_2\} + \sum_{v\in V(G_2)} \{d_{G_2}(v) + n_1\}\right] \end{aligned}$$

from where the desired result follows. □

Now we generalize the above result for n disjoint graphs $G_1, G_2, \ldots, G_n$.

Theorem 4. *Let $G_1, G_2, \ldots, G_n$ be n graphs with disjoint vertex sets $V_i = V(G_i)$ and edge sets $E_i = E(G_i), 1 \le i \le n$, without well-connected vertices. Then,*

$$C^{\xi}(G_1 + G_2 + ... + G_n) = \sum_{i=1}^{n} |E(G_i)| + \frac{1}{2}\left\{\sum_{i=1}^{n} |V(G_i)|\right\}^2 - \frac{1}{2}\sum_{i=1}^{n} |V(G_i)|^2.$$

Proof. From definition of join, we have $d_{G_1+G_2+...+G_n}(v_j) = d_{G_i}(v_j) + |V(G_1 + G_2 + ... + G_n)| - |V(G_i)|$ and since none of $G_1, G_2, ..., G_n$ contains any well connected vertex, we have $\varepsilon_{G_1+G_2+...+G_n}(v_j) = 2$ for all $v_j \in V(G_i)$.

Thus the connective eccentric index of $G_1 + G_2 + ... + G_n$ is given by

$$\begin{aligned} C^{\xi}(G_1 + G_2 + \ldots + G_n) &= \sum_{i=1}^{n}\sum_{j=1}^{|V(G_i)|} \frac{d_{G_1+G_2+...+G_n}(v_j)}{\varepsilon_{G_1+G_2+...+G_n}(v_j)} \\ &= \sum_{i=1}^{n}\sum_{j=1}^{|V(G_i)|} \frac{d_{G_i}(v_j) + |V(G_1 + G_2 + ... + G_n)| - |V(G_i)|}{2} \\ &= \frac{1}{2}\Big[|E(G_1)| + |E(G_2)| + ... + |E(G_n)| \\ &\quad + |V(G_1 + G_2 + ... + G_n)|\{|V(G_1)| + |V(G_2)| + ... \\ &\quad + |V(G_n)|\} - \left\{|V(G_1)|^2 + V(G_2)|^2 + ... + |V(G_n)|^2\right\}\Big] \end{aligned}$$

from where the desired result follows. □

The following corollaries are direct consequences of the theorem.

Corollary 1. *If nG denotes the join of n copies of G, then*

$$C^{\xi}(nG) = n|E(G)| + \frac{n(n-1)}{2}|V(G)|^2.$$



Figure 3: The complete n-partite graph.

Corollary 2. *Let $G = K_{m_1,m_2,...,m_n}$ be the complete n-partite graph having $v = |V(G)|$ number of vertices. Here the vertex set can be partitioned into n subsets $V_1, V_2, ..., V_n$ such that $V_1 \cup V_2 \cup ... \cup V_n = V(G)$ and $V_i \cap V_j = \phi, i \neq j, 1 \leq i, j \leq n$. From definition of join it is clear that $G = K_{m_1,m_2,...,m_n}$ is the join of n empty graphs with $m_1, m_2, ..., m_n$ number of vertices. Then the connective eccentric index of $G = K_{m_1,m_2,...,m_n}$ is given by*

$$C^{\xi}(K_{m_1,m_2,...,m_n}) = \sum_{\substack{1\leq i\leq n\\ 1\leq j\leq n\\ i\neq j}} m_i m_j.$$

2.3. Symmetric difference

Let G_1 and G_2 be two graphs with vertex sets $V(G_1)$ and $V(G_2)$ and edge sets $E(G_1)$ and $E(G_2)$. Then the symmetric difference of G_1 and G_2, denoted by $G_1 \oplus G_2$ is the graph with vertex set $V(G_1) \times V(G_2)$ in which any two vertices (u_1, u_2) is adjacent to (v_1, v_2) whenever u_1 is adjacent to v_1 in G_1 or u_2 is adjacent to v_2 in G_2, but not both. From definition of symmetric difference, the degree of a vertex (u_1, u_2) of $G_1 \oplus G_2$ is given by [13]

$$d_{G_1\oplus G_2}(u_1, u_2) = |V(G_2)|d_{G_1}(u_1) + |V(G_1)|d_{G_2}(u_2) - 2d_{G_1}(u_1)d_{G_2}(u_2).$$

Theorem 5. *The connective eccentric index of the symmetric difference $G_1 \oplus G_2$ of two graphs G_1 and G_2 is given by*

$$C^{\xi}(G_1 \oplus G_2) = |E(G_1)||V(G_2)|^2 + |E(G_2)||V(G_1)|^2 - 4|E(G_1)||E(G_2)|$$

where, none of G_1 and G_2 contains well-connected vertices.

Proof. Since the distance between any two vertices of a symmetric difference cannot exceed two, if none of the components contains well-connected vertices, the eccentricity of all vertices is constant and equal to two, i.e., $\varepsilon_{G_1\oplus G_2}(u_1, u_2) = 2$, for all vertices (u_1, u_2) [2, 9].

Thus the connective eccentric index of symmetric difference $G_1 \oplus G_2$ of two graphs G_1 and G_2 is given by

$$\begin{aligned} C^{\xi}(G_1 \oplus G_2) &= \sum_{(u_1,u_2)\in V(G_1\oplus G_2)} \frac{d_{G_1\oplus G_2}(u_1,u_2)}{\varepsilon_{G_1\oplus G_2}(u_1,u_2)} \\ &= \frac{1}{2}\sum_{u_1\in V(G_1)}\sum_{u_2\in V(G_2)} [|V(G_2)|d_{G_1}(u_1) + |V(G_1)|d_{G_2}(u_2) - 2d_{G_1}(u_1)d_{G_2}(u_2)] \\ &= \frac{1}{2}\Bigg[|V(G_2)|^2 \sum_{u_1\in V(G_1)} d_{G_1}(u_1) + |V(G_1)|^2 \sum_{u_2\in V(G_2)} d_{G_2}(u_2) \\ &\quad -2\sum_{u_1\in V(G_1)} d_{G_1}(u_1) \sum_{u_2\in V(G_2)} d_{G_2}(u_2)\Bigg] \end{aligned}$$

from where the desired result follows. □

2.4. Disjunction

The disjunction $G_1 \vee G_2$ of two graphs G_1 and G_2 is the graph with vertex set $V(G_1)\times V(G_2)$ in which (u_1, u_2) is adjacent to (v_1, v_2) whenever u_1 is adjacent with v_1 in G_1 or u_2 is adjacent with v_2 in G_2. Obviously, the degree of a vertex (u_1, u_2) of $G_1 \vee G_2$ is given by [2, 9]

$$d_{(G_1\vee G_2)}(u_1,u_2) = [|V(G_2)|d_{G_1}(u_1) + |V(G_1)|d_{G_2}(v_2) - d_{G_1}(u_1)d_{G_2}(v_2)]\,.$$

Theorem 6. *The connective eccentric index of the disjunction $G_1 \vee G_2$ of two graphs G_1 and G_2 is given by*

$$C^{\xi}(G_1 \vee G_2) = |E(G_1)||V(G_2)|^2 + |E(G_2)||V(G_1)|^2 - 2|E(G_1)||E(G_2)|$$

where none of G_1 and G_2 contains well-connected vertices.

Proof. Since the distance between any two vertices of a disjunction cannot exceed two, if none of the components contains well-connected vertices, the eccentricity of all vertices is constant and equal to two [2, 9]. Then the connective eccentric index of the disjunction $G_1 \vee G_2$ of two graphs G_1 and G_2 is computed as

$$\begin{aligned} C^{\xi}(G_1 \vee G_2) &= \sum_{(u_i,v_j)\in V(G_1\vee G_2)} \frac{d_{G_1\vee G_2}(u_1,u_2)}{\varepsilon_{G_1\vee G_2}(u_1,u_2)} \\ &= \frac{1}{2}\sum_{u_1\in V(G_1)}\sum_{v_2\in V(G_2)} [|V(G_2)|d_{G_1}(u_1) + |V(G_1)|d_{G_2}(u_2) - d_{G_1}(u_1)d_{G_2}(u_2)] \end{aligned}$$

from where the desired result follows. □

2.5. Splice

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are two graphs $V_1 \cap V_2 = \phi$. Let $v_1 \in V_1$ and $v_2 \in V_2$ be two given vertices of G_1 and G_1 respectively. A splice of G_1 and G_1 at the vertices v_1 and v_2 is obtained by identifying the vertices v_1 and v_2 in the union of G_1 and G_1 and is denoted by $S(G_1, G_2, v_1, v_2)$ [6]. Different topological indices of the splice graphs have already been computed [16]. Let $u \in V(S)$, then $\varepsilon_1(u)$ denotes the eccentricity of u as a vertex of G_1, $\varepsilon_2(u)$ denotes the eccentricity of u as a vertex of G_2 and $\varepsilon(u)$ denotes the eccentricity of u as a vertex of G. Let $d_S(x)$ be the degree of the vertex $x \in V(S)$. Then the connective eccentric index of $S(G_1, G_2, v_1, v_2)$ is computed as follows.

Theorem 7. *The connective eccentric index of splice of G_1 and G_2 is given by*

$$C^{\xi}(S) = \sum_{x \in V_1} \frac{d_S(x)}{max\{d_{G_1}(x, u_1) + \varepsilon(u_2)\varepsilon(x)\}} + \sum_{y \in V_2} \frac{d_S(y)}{max\{d_{G_2}(y, u_2) + \varepsilon_1(u_1)\varepsilon(y)\}}.$$

Proof. For any vertex $x \in V_1$, Sharafdini and Gutman [16] showed that

$$\varepsilon(x) = \max\{d_{G_1}(x, u_1) + \varepsilon_2(u_2)\varepsilon_1(x)\}.$$

Similarly, for any vertex $y \in V_2$,

$$\varepsilon(y) = \max\{d_{G_2}(y, u_2) + \varepsilon_1(u_1)\varepsilon(y)\}.$$

Thus, the desired result follows from the definition of connective eccentric index. □

References

[1] N. Alon, Eigenvalues and expanders, *Combinatorica* 6(1986) 83–96.

[2] A.R. Ashrafi, M. Ghorbani and M. A. Hossein-Zadeh, The eccentric connectivity polynomial of some graph operations, *Serdica J. Comput.* 5(2011) 101–116.

[3] P. Dankelmann, W. Goddard and C.S. Swart, The average eccentricity of a graph and its subgraphs, *Util. Math.* 65(2004) 41–51.

[4] N. De, Bounds for connective eccentric index, *Int. J. Contemp. Math. Sci.* 7(44)(2012) 2161–2166.

[5] T. Dehghan-Zadeh, H. Hua, A. R. Ashrafi and N. Habibi, Remarks on a conjecture about Randić index and graph radius, *Miskolc Math. Notes* 14(3)(2013) 845–850.

[6] T. Doslić, Splices, links and their degree-weighted Wiener polynomials, *Graph Th. Notes* 48(2005) 47–55.

[7] M. Ghorbani, Connective Eccentric Index of Fullerenes, *J. Math. Nanoscience* 1(2011) 43–52.

[8] M. Ghorbani and K. Malekjani, A new method for computing the eccentric connectivity index of fullerenes, *Serdica J. Comput.* 6(2012) 299–308.

[9] M. Ghorbani and M. A. Hosseinzadeh, A new version of Zagreb indices, *Filomat* 26(2012) 93–100.

[10] S. Gupta, M. Singh and A. K. Madan, Connective eccentricity Index: A novel topological descriptor for predicting biological activity, *J. Mol. Graph. Model.* 18(2000) 18–25.

[11] S. Hossein-Zadeh, A. Hamzeh and A. R. Ashrafi, Winer-type invariants of some graph operations, *Filomat* 23(3)(2009) 103–113.

[12] H. Hua, S. Zhang, K. Xu, Further results on the eccentric distance sum, *Disc. Appl. Math.* 160(2012) 170–180.

[13] W. Imrich and S. Klavžar, Product graphs: structure and recognition, John Wiley & Sons, New York, USA, 2000.

[14] M.H. Khalifeh, H. Yousefi-Azari and A. R. Ashrafi, The hyper-Wiener index of graph operations, *Comput. Math. Appl.* 56(5)(2008) 1402–1407.

[15] M.H. Khalifeh, H. Yousefi-Azari and A. R. Ashrafi, The first and second Zagreb indices of some graph operations, *Disc. Appl. Math.* 157(4)(2009) 804–811.

[16] R. Sharafdini and I. Gutman, Splice graphs and their topological indices, *Kragujevac J. Sci.* 35(2013) 89–98.

[17] G. Yu and L. Feng, On Connective Eccentricity Index of Graphs, *MATCH Commun. Math. Comput. Chem.* 69(2013) 611–628.